



Advanced Dermatology

Skin Cancer and
Laser Surgery Center, P.C.

www.advaderm.com

Board Certified Dermatologists

General Dermatology

Gregory G. Papadeas, D.O.
Steven A. Brenman, D.O.
Jamie G. Surovik, M.D.
Jennifer A. Ray, M.D.
Ryan G. Gamble, M.D.
Dawnielle C. Endly, D.O.
Ryan O'Leary, M.D.
Paul J. Grant, M.D.
Anne H. Hanson, D.O.
Roger I. Ceilley, M.D.

Mohs Micrographic Surgery

James T. Chapman, M.D.
Darlene S. Johnson, M.D.

Dermatopathology

Sean Amsbaugh, M.D.

Pediatric Dermatology

Elizabeth Swanson, M.D.



Medical Center of Aurora

1390 South Potomac Street
Suite 124
Aurora, CO 80012
303-368-8611
303-368-9791 fax

Centennial

12645 East Euclid Drive
Centennial, CO 80111
303-493-1910
303-493-1915 fax

Evergreen Medical Center

30960 Stagecoach Blvd., Suite W140
Evergreen, CO 80439
303-670-7065
303-670-7067 fax

Summit County

710 Summit Blvd., Suite 102
PO Box 4005
Frisco, CO 80443
970-668-9650
970-668-9654 fax

Castle Rock

2352 Meadows Blvd., Suite 220
Castle Rock, CO 80109
303-493-1910

JUVEDERM VOLUMA XC PRE & POST-OP CARE

Pre Treatment Instructions

- To reduce the risks of bruising and/or bleeding, please avoid taking the following one week prior to your treatment and one week following your treatment.
 - Aspirin
 - St. John's Wort
 - High doses of Vitamin E
 - Anti-inflammatory medications
 - Ibuprofen
 - Fish Oil
- Avoid consumption of alcohol within 24 hours of the procedure and for 48 hours after the procedure.

Post Treatment Instructions

- Avoid touching the treated area within 6 hours following treatment. After that, the area can be gently washed with soap and water.
- Until the initial redness and swelling have resolved, avoid exposure of the treated area to intense heat (sunlamp or sunbathing).
- Immediately after the treatment, the most commonly reported side effects were temporary redness, swelling and/or bruising at the injection site. These effects typically resolve within a few days. Cold compresses may be used immediately after treatment for 5-10 minutes every 2-3 hours to reduce the chance of swelling and/or bruising.
- If you have previously suffered from facial cold sores, there is a risk that the needle punctures could contribute to another recurrence. Speak to your physician about medications that may minimize a recurrence.
- Avoid exercise, spicy foods and alcohol for 6 hours post-treatment.

If you have any questions, please don't hesitate to call the office.