



Advanced[®] Dermatology

Skin Cancer and
Laser Surgery Center, P.C.

www.advaderm.com

Board Certified Dermatologists

General Dermatology

Gregory G. Papadeas, D.O.
James T. Chapman, M.D.
Steven A. Brenman, D.O.
Jamie G. Surovik, M.D.
Jennifer A. Ray, M.D.
Ryan G. Gamble, M.D.
Dawnielle C. Endly, D.O.
Ryan O'Leary, M.D.
Powell Perng, M.D.
Anne H. Hanson, D.O.
Roger I. Ceilley, M.D.

Mohs Micrographic Surgery

Darlene Johnson Skow, M.D.
Misha D. Miller, M.D.
Jamison E. Strahan, M.D.

Dermatopathology

Sean Amsbaugh, M.D.

Pediatric Dermatology



Medical Center of Aurora

1390 South Potomac Street
Suite 124
Aurora, CO 80012
303-368-8611
303-368-9791 fax

Centennial

12645 East Euclid Drive
Centennial, CO 80111
303-493-1910
303-493-1915 fax

Evergreen Medical Center

30960 Stagecoach Blvd., Suite W140
Evergreen, CO 80439
303-670-7065
303-674-6055 fax

Summit County

710 Summit Blvd., Suite 102
PO Box 4005
Frisco, CO 80443
970-668-9650
970-668-9654 fax

Castle Rock

2352 Meadows Blvd., Suite 225
Castle Rock, CO 80109
303-493-1910

EFUDEX

Efudex is an anti-cancer medicine that has been made into a cream. It is used to treat the pre-cancerous skin lesions called actinic keratoses. It is also sometimes used to treat very thin skin cancers. It does not work on thicker cancers or benign lesions, such as seborrheic keratoses or milia.

Usually, Efudex has no effect on normal skin. If you were to apply it to an area of skin that is almost never exposed to sunlight, such as an under-arm, nothing would happen. When applied to skin that has previously been sun-exposed, it leaves the non-cancerous skin alone, but gradually kills the pre-cancerous spots. When we use Efudex, we treat an area of skin; we do not spot treat. To try to spot treat takes away the great advantage of Efudex; it kills all the pre-cancers, even the ones that you or your doctor cannot see. This, along with careful use of sunscreen, prevents you from getting as many (or sometimes any!) new actinic keratoses in the future. Another advantage of Efudex is that it has no effect on the lower layers of the skin. For that reason, it usually doesn't cause any scarring.

The biggest disadvantage of Efudex is the same as its biggest advantage: it kills all the pre-cancers. This means, if you have a lot of them, that you will end your Efudex treatment with a lot of small erosions, which can itch, sting, or burn. With proper care, these heal rapidly, but there is usually a weeklong period of discomfort. We can minimize this discomfort, but we cannot eliminate it. The second disadvantage is that at the peak of the reaction, the treated area is unsightly. The drying pre-cancers get very red, then weep and crust. Most people prefer to be out of the public eye during this time. Even after the spots are completely healed over, they may remain pink for an extended period of time, sometimes a few months.

Efudex produces a temporary but severe sun sensitivity. During the treatment period, which usually lasts 2 to 3 weeks, the treated skin must be protected from sunlight and even the scattered light from a blue sky. The safest policy is to stay indoors while the sun is up. The days are much shorter in the months of November through January, so many treat their patients during this time. If you must go outside during daylight, you must apply a thick coat of SPF 30 or greater and limit your time in the sun to 10 minutes or less. Some areas, such as the scalp, arms, hands, can be covered with a hat, long sleeves, and gloves. If the face is not being treated, and the areas being treated are covered, you may then go outside for indefinite periods. Some people are, or become, allergic to Efudex. In these individuals it produces an itchy rash, quite unlike the anti-cancer reaction, all over the treated area. These people cannot use Efudex.

Many people have large numbers of pre-cancers. There can be so many of them that they grow together, and it becomes impossible to tell where the normal skin ends, and the pre-cancerous skin begins. In this situation it is impractical to freeze each of these pre-cancers. If you are reading this, it is because your physician thinks that in your case the advantages of using Efudex outweigh the disadvantages. If you would like to use Efudex, please inform your doctor, and we will set up a time to begin treatment.



Advanced[®] Dermatology

Skin Cancer and
Laser Surgery Center, P.C.

www.advaderm.com

Board Certified Dermatologists

General Dermatology

Gregory G. Papadeas, D.O.
James T. Chapman, M.D.
Steven A. Brenman, D.O.
Jamie G. Surovik, M.D.
Jennifer A. Ray, M.D.
Ryan G. Gamble, M.D.
Dawnielle C. Endly, D.O.
Ryan O'Leary, M.D.
Powell Perng, M.D.
Anne H. Hanson, D.O.
Roger I. Ceilley, M.D.

Mohs Micrographic Surgery

Darlene Johnson Skow, M.D.
Misha D. Miller, M.D.
Jamison E. Strahan, M.D.

Dermatopathology

Sean Amsbaugh, M.D.

Pediatric Dermatology



Medical Center of Aurora

1390 South Potomac Street
Suite 124
Aurora, CO 80012
303-368-8611
303-368-9791 fax

Centennial

12645 East Euclid Drive
Centennial, CO 80111
303-493-1910
303-493-1915 fax

Evergreen Medical Center

30960 Stagecoach Blvd., Suite W140
Evergreen, CO 80439
303-670-7065
303-674-6055 fax

Summit County

710 Summit Blvd., Suite 102
PO Box 4005
Frisco, CO 80443
970-668-9650
970-668-9654 fax

Castle Rock

2352 Meadows Blvd., Suite 225
Castle Rock, CO 80109
303-493-1910

Efudex / Carac/ Solaraz(Fluorouracil)

1. You have been prescribed a topical medication for the treatment of pre-cancerous and early cancer lesions. It should be used as directed for the complete duration of the time recommended. This is very important to maximize the effectiveness of the treatment.

****If you get cold sores, or have a history of Herpes/Cold sores, please notify you doctor. You will need to take another medication to prevent cold sores from starting during your Efudex treatment.**

2. Prior to applying the topical medication, wash your face with mild cleanser (e.g., Dove[®]/Cetaphil[®]) and pat it dry. There is no need to scrub the area vigorously, as this will only irritate the skin.
3. Allow the skin to dry. Applying the medication to wet skin will increase irritation.
4. Apply a thin layer of the medication and lightly rub it in.
5. If you use moisturizer / sunscreen, apply it 30 minutes after the medication has been applied. Generally, sun exposure will make you more uncomfortable.
6. Repeat this application process as often as is recommended on your prescription.
7. It is normal that most people using these medications will experience redness and irritation for the skin in the areas treated (may look like poison ivy). This is an indication that the medication is effective. In general, the worse the irritation – the better the results. For some people it may take 2 weeks or longer. Please DO NOT STOP your medication thinking that it is not working for you!
8. We recommend applying a light face moisturizer (Cetaphil[®]/ Oil of Olay[®] etc.) or Aquaphor[®] (this is thick Vaseline[®] type moisturizer) as often as needed to keep the skin and/or scale soft. If this occurs, the medication will not be able to penetrate the scab and treatment will not be as effective as necessary.
9. If any areas become open or weepy, we encourage you to apply Bacitracin antibiotic ointment at least twice a day to prevent infection. Then apply a thick coat of Aquaphor[®].
10. If your skin becomes itchy, you may apply over-the-counter Hydrocortisone[®] 1% ointment/cream up to twice daily until the itch sensation stops.
11. If you have further questions or concerns, please call our office rather than stopping the medication without contacting us.
12. **Keep out of reach to children and pets.**

Follow -up Care.

1. Return to our office in _____ for follow-up care.
2. Have a complete skin check in 6 months from your biopsy date, and every 6 months after that for 2 years.

IF YOU HAVE ANY QUESTIONS OR CONCERNS, PLEASE CALL OUR OFFICE.